

  
**MYOBLOC<sup>®</sup>**  
rimabotulinumtoxinB  
Injection [5,000 Units/mL]

Not actual healthcare providers.



# CIRCLE OF CARE<sup>™</sup> REFERENCE GUIDE

ACCESS AND USE THE CIRCLE OF CARE PROVIDER PORTAL  
FOR EASY AND CONVENIENT MYOBLOC REIMBURSEMENT

## INDICATIONS

MYOBLOC<sup>®</sup> injection is indicated for:

- the treatment of cervical dystonia (CD) to reduce the severity of abnormal head position and neck pain associated with CD in adults
- the treatment of chronic sialorrhea in adults

## IMPORTANT SAFETY INFORMATION

### **WARNING: DISTANT SPREAD OF TOXIN EFFECT**

See full prescribing information for complete boxed **WARNING**.


The effects of MYOBLOC<sup>®</sup> and all botulinum toxin products may spread from the area of injection to produce symptoms consistent with botulinum toxin effects. These symptoms have been reported hours to weeks after injection. Swallowing and breathing difficulties can be life threatening and there have been reports of death. The risk of symptoms is probably greatest in children treated for spasticity but symptoms can also occur in adults, particularly in those patients who have an underlying condition that would predispose them to these symptoms.

Please refer to the full [Prescribing Information](#), including [Boxed WARNING](#), and additional Important Safety Information throughout, and on pages [4](#) and [5](#).

## WHAT IS THE CIRCLE OF CARE?

MYOBLOC comes with robust access and a suite of support services to assist your practice and to help appropriate patients get the treatment they need. **The Circle of Care Provider Portal** is an integral part of the MYOBLOC Support Services Suite.

The Circle of Care Provider Portal allows for online submissions of insurance benefit investigations and prior authorization facilitation for new and current MYOBLOC patients.



## CIRCLE OF CARE GIVES YOU EASY ACCESS TO:

- Prior authorization for existing and new patients
- Specialty pharmacy communication and Patient Assistance Programs
- Online submission for insurance benefit investigations including 2-business day turnaround in most cases
- Real-time patient status updates and ability to track your patient's journey
- Access important forms and begin the Patient Referral process

### IMPORTANT SAFETY INFORMATION (cont'd)

#### CONTRAINDICATIONS

MYOBLOC is contraindicated in patients with:

- A known hypersensitivity to any botulinum toxin product or to any of the components in the formulation
- Infection at the proposed injection site(s)

**Please refer to the full [Prescribing Information](#), including [Boxed WARNING](#), and additional Important Safety Information throughout, and on [pages 4](#) and [5](#).**

# GETTING STARTED WITH CIRCLE OF CARE

---

## Step 1

Visit <https://hcp.supernusconnect.com/#/login>

## Step 2

Enter your credentials (new users will be guided through a registration for your practice)

## Step 3

After entering your information, you will have the option to add additional practice addresses and additional providers on the next screen

## Step 4

Once your account is approved, you will receive a registration approval email and gain access to your dashboard for an overview of patients and claims status



Scan the code to visit the Circle of Care  
Provider Portal and get started today.

## IMPORTANT SAFETY INFORMATION (cont'd)

### WARNINGS AND PRECAUTIONS

#### ▪ Lack of Interchangeability Between Botulinum Toxin Products

The potency units of MYOBLOC are specific to the preparation and biological activity assay method utilized. Due to differences in the aspects of this assay such as the vehicle, dilution scheme, and laboratory protocols for various potency assays, potency units are not interchangeable with other preparations of botulinum toxin products and, therefore, units of biological activity of MYOBLOC cannot be compared to or converted into units of any other botulinum toxin products assessed with any other specific assay method.

Please refer to the full [Prescribing Information](#), including [Boxed WARNING](#), and additional Important Safety Information throughout, and on pages [4](#) and [5](#).

## BILLING INFORMATION

Condition	Code	Description
Cervical dystonia*	ICD10 - G24.3	Spasmodic torticollis (CD)
	CPT - 64616	Chemodeneration of muscle(s), cervical spinal muscle(s); 50 modifier for bilateral injections
	J0587	MYOBLOC is billed in 100 units <i>(ie, 5,000 units = 50 units billed)</i>
	95873	Electrical stimulation for guidance in conjunction with chemodeneration
	95874	Needle electromyography for guidance in conjunction with chemodeneration
	76942	Ultrasonic guidance for needle placement imaging
Chronic sialorrhea*	ICD10 - K11.7	Disturbances of salivary secretion (sialorrhea)
	CPT - 64611	Chemodeneration of parotid and submandibular salivary glands, bilateral
	J0587	MYOBLOC is billed in 100 units <i>(ie, 5,000 units = 50 units billed)</i>
	76942	Ultrasonic guidance for needle placement imaging

\*Table contains possible appropriate codes for cervical dystonia and chronic sialorrhea.

For more information on reimbursement for MYOBLOC, including billing and coding, visit [www.MYOBLOC.com](http://www.MYOBLOC.com).

### IMPORTANT SAFETY INFORMATION (cont'd)

#### WARNINGS AND PRECAUTIONS (cont'd)

##### ▪ Hypersensitivity Reactions

Serious hypersensitivity reactions have been reported with botulinum toxin products. Angioedema, urticaria, and rash have occurred with MYOBLOC treatment. Hypersensitivity reactions can also include anaphylaxis, serum sickness, soft tissue edema, and dyspnea. If serious and/or immediate hypersensitivity reactions occur, discontinue further injection of MYOBLOC and institute appropriate medical therapy immediately. The use of MYOBLOC in patients with a known hypersensitivity to any botulinum neurotoxin or to any of the excipients (human albumin, sucrose), could lead to a life-threatening allergic reaction.

Please refer to the full [Prescribing Information](#), including [Boxed WARNING](#), and additional Important Safety Information throughout, and on pages [4](#) and [5](#).

## IMPORTANT SAFETY INFORMATION

### **WARNING: DISTANT SPREAD OF TOXIN EFFECT**

See full prescribing information for complete boxed WARNING.

The effects of MYOBLOC® and all botulinum toxin products may spread from the area of injection to produce symptoms consistent with botulinum toxin effects. These symptoms have been reported hours to weeks after injection. Swallowing and breathing difficulties can be life threatening and there have been reports of death. The risk of symptoms is probably greatest in children treated for spasticity but symptoms can also occur in adults, particularly in those patients who have an underlying condition that would predispose them to these symptoms.

### **CONTRAINDICATIONS**

MYOBLOC is contraindicated in patients with:

- A known hypersensitivity to any botulinum toxin product or to any of the components in the formulation
- Infection at the proposed injection site(s)

### **WARNINGS AND PRECAUTIONS**

#### **▪ Lack of Interchangeability Between Botulinum Toxin Products**

The potency units of MYOBLOC are specific to the preparation and biological activity assay method utilized. Due to differences in the aspects of this assay such as the vehicle, dilution scheme, and laboratory protocols for various potency assays, potency units are not interchangeable with other preparations of botulinum toxin products and, therefore, units of biological activity of MYOBLOC cannot be compared to or converted into units of any other botulinum toxin products assessed with any other specific assay method.

#### **▪ Hypersensitivity Reactions**

Serious hypersensitivity reactions have been reported with botulinum toxin products. Angioedema, urticaria, and rash have occurred with MYOBLOC treatment. Hypersensitivity reactions can also include anaphylaxis, serum sickness, soft tissue edema, and dyspnea. If serious and/or immediate hypersensitivity reactions occur, discontinue further injection of MYOBLOC and institute appropriate medical therapy immediately. The use of MYOBLOC in patients with a known hypersensitivity to any botulinum neurotoxin or to any of the excipients (human albumin, sucrose), could lead to a life-threatening allergic reaction.

#### **▪ Dysphagia and Breathing Difficulties**

Treatment with MYOBLOC and other botulinum toxin products can result in swallowing or breathing difficulties. Patients with pre-existing swallowing or breathing difficulties may be more susceptible to these complications. In most cases, this is a consequence of weakening of muscles in the area of injection that are involved in breathing or swallowing. When distant effects occur, additional respiratory muscles may be involved. Deaths as a complication of severe dysphagia have been reported after treatment with botulinum toxin. Dysphagia may persist for several months and require use of a feeding tube to maintain adequate nutrition and hydration. Aspiration may result from severe dysphagia and is a particular risk when treating patients in whom swallowing or respiratory function is already compromised.

**Please refer to the full [Prescribing Information](#), including [Boxed WARNING](#), and additional [Important Safety Information](#) on [page 5](#).**

## WARNINGS AND PRECAUTIONS (cont'd)

### ▪ Dysphagia and Breathing Difficulties (cont'd)

- **Cervical Dystonia:** Treatment of cervical dystonia with botulinum toxins may weaken neck muscles that serve as accessory muscles of ventilation. This may result in a critical loss of breathing capacity in patients with respiratory disorders who may have become dependent upon these accessory muscles. There have been postmarketing reports of serious breathing difficulties, including respiratory failure, in cervical dystonia patients. Patients treated with botulinum toxin may require immediate medical attention should they develop problems with swallowing, speech or respiratory disorders. These reactions can occur within hours to weeks after injection with botulinum toxin.
- **Pre-Existing Neuromuscular Disorders:** Individuals with peripheral motor neuropathic diseases, amyotrophic lateral sclerosis, or neuromuscular junctional disorders (e.g., myasthenia gravis or Lambert-Eaton syndrome) should be monitored particularly closely when given botulinum toxin. Patients with neuromuscular disorders may be at increased risk of clinically significant effects including severe dysphagia and respiratory compromise from typical doses of MYOBLOC.

### ▪ Human Albumin and Transmission of Viral Diseases

This product contains albumin, a derivative of human blood. Based on effective donor screening and product manufacturing processes, it carries an extremely remote risk for transmission of viral diseases and variant Creutzfeldt-Jakob disease (vCJD). There is a theoretical risk for transmission of Creutzfeldt-Jakob disease (CJD), but if that risk actually exists, the risk of transmission would be considered extremely remote. No cases of transmission of viral diseases, CJD, or vCJD have ever been identified for licensed albumin or albumin contained in other licensed products.

## MOST COMMON ADVERSE REACTIONS (>5% of patients and >5% more than placebo)

**Cervical Dystonia:** dry mouth, dysphagia, injection site pain, headache

**Chronic Sialorrhea:** dry mouth, dysphagia

## DRUG INTERACTIONS

**Aminoglycosides and Other Agents Interfering with Neuromuscular Transmission:** Co-administration of MYOBLOC and aminoglycosides or other agents interfering with neuromuscular transmission (e.g., curare-like compounds) should only be performed with caution as the effect of the toxin may be potentiated. **Anticholinergic Drugs:** Use of anticholinergic drugs after administration of MYOBLOC may potentiate systemic anticholinergic effects. **Other Botulinum Neurotoxin Products:** The effect of administering different botulinum toxin products at the same time or within several months of each other is unknown. Excessive neuromuscular weakness may be exacerbated by administration of another botulinum toxin prior to the resolution of the effects of a previously administered botulinum toxin. **Muscle Relaxants:** Excessive weakness may also be exaggerated by administration of a muscle relaxant before or after administration of MYOBLOC.

## INDICATIONS

MYOBLOC® injection is indicated for:

- the treatment of cervical dystonia (CD) to reduce the severity of abnormal head position and neck pain associated with CD in adults
- the treatment of chronic sialorrhea in adults

Please refer to the full [Prescribing Information](#), including **Boxed WARNING**.

# MY SUPPORT. MY PATIENTS. MYOBLOC.

Not actual healthcare provider.



DO YOUR NEXT REIMBURSEMENT ON  
THE CIRCLE OF CARE PROVIDER PORTAL.  
VISIT [MYOBLOCHCP.COM/REIMBURSEMENT](https://myoblochcp.com/reimbursement).

Print page [7](#) for quick reference to keep in your practice

## INDICATIONS

MYOBLOC® injection is indicated for:

- the treatment of cervical dystonia (CD) to reduce the severity of abnormal head position and neck pain associated with CD in adults
- the treatment of chronic sialorrhea in adults

## IMPORTANT SAFETY INFORMATION


### WARNING: DISTANT SPREAD OF TOXIN EFFECT

See full prescribing information for complete boxed WARNING.

The effects of MYOBLOC® and all botulinum toxin products may spread from the area of injection to produce symptoms consistent with botulinum toxin effects. These symptoms have been reported hours to weeks after injection. Swallowing and breathing difficulties can be life threatening and there have been reports of death. The risk of symptoms is probably greatest in children treated for spasticity but symptoms can also occur in adults, particularly in those patients who have an underlying condition that would predispose them to these symptoms.

Please refer to the full [Prescribing Information](#), including [Boxed WARNING](#), and additional Important Safety Information throughout and on pages [4](#) and [5](#).

MYOBLOC is a registered trademark of Solstice Neurosciences, LLC, a wholly-owned subsidiary of Supernus Pharmaceuticals, Inc. Circle of Care is a trademark of Solstice Neurosciences, LLC. © 2023. Supernus Pharmaceuticals, Inc. MYO.2023-0024 05/23.





## FOR EASY REIMBURSEMENT FOR YOUR SUPERNUS INJECTIONS

---

### Step 1

Visit <https://hcp.supernusconnect.com/#/login>

### Step 2

Enter your credentials (new users will be guided through a registration for your practice)

### Step 3

After entering your information, you will have the option to add additional practice addresses and additional providers on the next screen

### Step 4

Once your account is approved, you will receive a registration approval email and gain access to your dashboard for an overview of patients and claims status



Scan the code to visit the Circle of Care  
Provider Portal and get started today.